



***Calpurnia aurea* (Alt.) Benth: A Review of Its Ethnobotanical, Phytochemical, and Pharmacological Properties**

Asaye Asfaw ^{1*} and Wondimagegn Mengist ¹

¹ Department of natural resource management, Debre Birhan University, Ethiopia

Abstract

(*Calpurnia aurea* (Alt.) Benth is an accepted name for a species within the genus *Calpurnia*, belonging to the family Fabaceae. The primary objective of this review was to explore the traditional uses of *Calpurnia aurea* (*C. aurea*) for the treatment of various human and livestock diseases. The study evaluated utilization of parts of this plant for mediation, its methods of preparation, and routes of administration. It included studies on phytochemical extraction, characterization, and pharmacological activities. However, as the main objective of this study was to review studies on phytochemical extraction, characterization, and pharmacological activities, the word “also” should be avoided. The review also includes findings related to phytochemical extraction, characterization, and the pharmacological activities associated with *C. aurea*. It relied on secondary information sourced from relevant ethnobotanical literatures and published studies. Although practices regarding its use may vary, local populations commonly employ the roots, leaves, and seeds of the plant to treat various ailments in humans and livestock, because it’s notable bioactive compounds. *C. aurea* contains a diverse array of secondary metabolites, including alkaloids, terpenoids, saponins, cardiac glycosides, steroids, phenols, flavonoids, anthraquinones, and tannins. While various pharmacological properties of the extracts have been experimentally tested, comprehensive scientific explorations and detailed investigations are still needed.

Keywords: Antimicrobial activities, Lattice; *Calpurnia aurea*; phytochemical analysis, traditional uses

1. Introduction

Humans to treat and manage diseases since ancient times have utilized medicinal plants [1]. Several studies indicate that medicinal plants are still in use in both developed and developing countries, largely due to the emergence of drug-resistant bacteria and the increasing adverse effects associated with chemical medications [1, 2]. The plant family Fabaceae is one of the largest Family in the number of medicinal plants [1]. The family Fabaceae encompasses all legumes, commonly referred to as pea or bean plants. Among the largest families of flowering plants worldwide, Fabaceae ranks as the third largest, comprising approximately 20,000 species classified into about 800 genera [3]. The most prevalent and largest genera include *Astragalus*, which contains over 2,400 species; *Acacia*, with more than 950 species; and *Indigofera* and *Crotalaria*, each with around 700 species. Additionally, *Mimosa* has approximately 500 species, collectively representing about 9.4% of all flowering

plant species [4]. The family Leguminosae is further divided into three subfamilies: Papilionoideae, Caesalpinioideae, and Mimosoideae. These subfamilies are sometimes recognized as separate and independent families, and their flowers are useful for identification.

They exhibit extremely diverse characteristics [5]. The primary and distinctive feature of this family is the presence of root nodules that contain *Rhizobium* bacteria, which are capable of converting atmospheric nitrogen (inert) into usable forms. Species within this family range from herbs to trees [6]. Occasionally, these subfamilies are recognized as separate and independent families.

Economically, they represent the third largest group of plant families and are utilized as crops, green manures, pulses, oil crops, fruits, root and tuber crops, industrial crops, and forage species [7]. Additionally, they are

*Corresponding author: leulasaye@yahoo.com

Received: Dec. 20/2024, Revised Nov. 10/2024, Accepted: Jan 17/2025, Published: Feb 1/2025.

Asaye Asfaw and Wondimagegn Mengist

essential for synthesizing a wide variety of natural products, including flavors, poisons, dyes, and hold significant importance for medicinal purposes [8].

The members of this family are cosmopolitan in their distribution and abundant in forests [7]. Traditionally, many Fabaceae plants have provided safe and effective therapies for the treatment of various diseases in both humans and livestock [1].

C. aurea belongs to the Fabaceae family and is widely distributed in India, as well as in Africa, particularly in South Africa and eastern Africa, especially in Ethiopia and Eritrea [9]. This species thrives in fertile, well-drained soil with ample water during the summer [10]. It typically grows in forests, along forest margins, and in clearings. In forested areas, it can reach heights of 9 to 15 meters, while in open areas, it is more commonly found as a shrub or small tree, measuring 3 to 4 meters tall [11].

C. aurea has numerous traditional medicinal applications. Local communities utilize its leaves and seeds to treat various internal and external ailments in both humans and livestock. Several studies have demonstrated that *C. aurea* is effective in preventing insect bites and ectoparasites in livestock [12]. A variety of phytochemical compounds with medicinal properties have been isolated from different parts of the plant by various researchers [13-15]. These compounds exhibit antimicrobial and antifungal effects [16, 17]. However, many of these studies remain at a preliminary stage, indicating a need for more comprehensive investigations, particularly regarding phytochemical extraction, isolation, and pharmacological testing.

Nowadays, the study of how local communities utilize medicinal plants is a major focus in the field of ethnobotany. As a result, numerous research projects on ethnobotanical uses, phytochemical analysis, and efficacy testing are being conducted in various parts of the world. However, uses of specific medicinal plants, their bioactive compounds, and their roles in modern applications have not been compiled separately.

Therefore, the primary purpose of this paper was to review the status of *C. aurea* in terms of its traditional medicinal uses, phytochemical extractions, and pharmacological activities.

2. Review of Methods

To achieve the objectives of this review a comprehensive examination of books, review articles, original research journal publications, and ethnobotanical texts on *C. aurea* was conducted. A literature search for this review involved an online search of established databases to access books and reputable scholarly articles. The search was limited to articles written in English and was conducted between March 1900 and April 2024, using key search terms

such as “ethnobotany of *C. aurea*,” “traditional uses of *C. aurea*,” “phytochemical profile of *C. aurea*,” “active ingredients of *C. aurea*,” and “diseases treated by *C. aurea*,” were among others.

This study searched Clarivate's Web of Science, Scopus, Google Scholar, and PubMed for papers published from 1990 to April 2024 with the topics “ethnobotany of *C. aurea*,” “traditional uses of *C. aurea*,” “phytochemical profile of *C. aurea*,” “active ingredients of *C. aurea*,” and “diseases treated by *C. aurea*”. The earliest related literature found was published in 1984, so the researchers adjusted the time-span of the literature search from 1984 to 2024. The criteria for including articles in this review were: (i) articles must be published in English language and (ii) literature must be from peer-reviewed scientific journals and institutional documents published until April 2024. However, this review work excluded non English articles

3. Results and Discussion

3.1 A Botanical Description of *Calpurnia aurea*

The genus *Calpurnia* comprises several species, some of which are endemic to South Africa. *C. aurea* is the most widespread species, commonly found in various regions of sub-Saharan Africa and present in southern India. The name of the genus is derived from Calpurnius, a Roman poet who sought to emulate the renowned Virgil. consequently, the name Calpurnius was selected for the morphologically similar genus *Calpurnia* [18]

C. aurea is a small, multi-stemmed tree, typically reaching heights of 1 to 5 meters (occasionally up to 10 meters) but rarely exceeding 5 meters. The species features a stem diameter of up to 10 centimeters, with bark that is pale brown and darkens with age. The leaves are arranged alternately in a compound formation on stalks approximately 25 centimeters long, consisting of 5 to 15 pairs of leaflets plus a terminal leaflet. These leaflets are oblong, pale green, and measure about 2.5 centimeters, each ending in a fine hair-like tip. The flowers are bright yellow and pea-shaped, measuring around 2.5 centimeters across, and are clustered in dense heads on stalks that range from 7 to 24 centimeters in length. The largest petal appears almost split in half (Figure 1.A). The fruit consists of thin, membranous pods measuring 12 centimeters in length, pale yellow-brown in color, with one edge slightly winged, containing 4 to 8 seeds. The unopened pods (Figure 1.B) remain on the tree for an extended period.

Asaye Asfaw and Wondimagegn Mengist



Figure 1. A. The golden yellow flower of *C. aurea* B. Pod and seeds of *C. aurea*

Class	Magnoliopsida
Order	Fabales
Family	Fabaceae
Genus	Calpurnia
Species	<i>Calpurnia. aurea</i>

3.2 ECOLOGY AND GEOGRAPHIC DISTRIBUTION OF CALPURNIA AUREA

C. aurea thrives in the highlands of East Africa, extending from Eritrea in the north to as far west as the Central African Republic, Zaire, and Angola, and reaching south to South Africa. It is also prevalent in southern India [19]. In Ethiopia, it is particularly abundant in the mountainous regions across all provinces [19]. This species is commonly found in forest margins, as well as in bushland and grassland areas that are favored by overgrazing. The altitudinal range of *C. aurea* spans from 1,650 to 2,550 meters (up to 3,000 meters) above sea level throughout Ethiopia [19]. The geographical distribution of *C. aurea* in Ethiopia has been mapped using coordinates from ethnobotanical research papers, with the most common occurrences found in the middle and highland areas of the country as indicated in Figure 2.




Figure 1. The geographical distribution of *C. aurea* species in Ethiopia

3.3 PROPAGATION METHODS

C. aurea utilizes its flowers as reproductive organs and propagates through its seeds and cuttings. The plant blooms nearly year-round, with the majority of flowers appearing after the rainy season. Calpurnia is easy to cultivate.

3.4 MEDICINAL AND OTHER USES OF CALPURNIA AUREA

Although the local community utilizes the species in various ways, the most common applications include

Source...accessed from [http:// www.truthfinder.com](http://www.truthfinder.com) on 26 February 2025

As shown in Table 1, the species has different names in Ethiopia across different localities and societies.

Table 1. Local names of *C. aurea* in Ethiopia

Local names	
Amharic	Digita
Afan Oromo	Ceekaa
Tgrigna	Hetsawus
English name	Natal Laburnum
Sidamigna	Chekata

C. aurea is a multi-stemmed woody shrub plant that belongs to the Fabaceae family and Table 2 shows its detailed botanical classification.
Table 2. Scientific classification of *C. aurea*

Taxonomic classification	
Kingdom	Plantae
Division	Magnoliophyta

Asaye Asfaw and Wondimagegn Mengist

firewood, agricultural tools, bee forage, fish poison for harvesting, lice protection, live fencing, walking sticks, and ornamental purposes. Additionally, the locals use extracts from the leaves and seeds to treat various ailments, employing them as anti-rabies, anti-diarrheal, and anti-malaria agents [16]. Many research studies have indicated that *C. aurea* has various traditional medicinal properties [20]. Ethnobotanical studies have established that *C. aurea* is highly valued as a traditional medicinal plant. The findings suggest that different parts of the plant are utilized to treat a range of human and livestock ailments. This indicates that locally, people use it to treat various diseases affecting both humans and livestock. Moreover, it shows different ethnic groups, both in Ethiopia and beyond, utilize various parts of *C. aurea* to address a wide range of ailments

Traditional used parts of <i>C.urena</i>	Locally used to treated	Reported by
Leaf /Root/ seeds/fruits/stem barks/	Snake bite	T Teklehaymanot and M Giday [21]; M Megersa, Z Asfaw, E Kelbessa, A Beyene and B Woldeab [22]; S Zerabruk and G Yirga [23].
Leaf	Stomach-ache	B Kidane, T van Anandel, LJG van der Maesen and Z Asfaw [24]; T Teklehaymanot [25]; S Zerabruk and G Yirga [23]; H Tadege, E Mohammed, K Asres and T Gebre-Mariam [26].
Leaf /Root/ seeds/fruits/stem barks/	Diarrhea	AA Adedapo, FO Jimoh, S Koduru, AJ Afolayan and PJ Masika [9]; S Umer, A Tekewe and N Kebede [27]; G Alemayehu, Z Asfaw and E Kelbessa [28]; R Regassa, T Bekele and M Megersa [29].
Leaf	Malaria	K Tolossa, E Debela, S Athanasiadou, A Tolera, G Ganga and JG Houdijk [30]; ; M Waka, RJ Hopkins and C Curtis [31].

Leaf /Root/ seeds/fruits/stem barks	Amoebic dysentery	E d'Avigdor, H Wohlmuth, Z Asfaw and T Awas [32]; T Teklehaymanot [25]; B Adugna, NK GetachewTerefe, W Mamo and S Keskes [33].
Leaf	Rabies	H Yineger, D Yewhalaw and D Teketay [34]; H Tadege, E Mohammed, K Asres and T Gebre-Mariam [26]; M Giday, T Teklehaymanot, A Animut and Y Mekonnen [35]; A Asfaw, E Lulekal, T Bekele, A Debella, A Abebe and S Degu [1]
Leaf /Root/ seeds/fruits/stem barks	Diabetes mellitus	M Giday, T Teklehaymanot, A Animut and Y Mekonnen [35].
Leaf	Skin infections	K Tolossa, E Debela, S Athanasiadou, A Tolera, G Ganga and JG Houdijk [30]; T Teklehaymanot [25]; H Agisho, M Osie and T Lambore [36].
Leaf	Swelling or cancer	H Tadege, E Mohammed, K Asres and T Gebre-Mariam [26].
Leaf	Syphilis	H Tadege, E Mohammed, K Asres and T Gebre-Mariam [26]; AA Adedapo, FO Jimoh, S Koduru, AJ Afolayan and PJ Masika [9]
Leaf	Leishmaniasis	AA Adedapo, FO Jimoh, S Koduru, AJ Afolayan and PJ Masika [9]; H Tadege, E Mohammed, K Asres and T Gebre-Mariam [26]
Leaf	Tuberculous	H Tadege, E

Asaye Asfaw and Wondimagegn Mengist

	osis	Mohammed, K Asres and T Gebre-Mariam [26];
Leaf	Elephantiasis	AA Adedapo, FO Jimoh, S Koduru, AJ Afolayan and PJ Masika [9]
Leaf /Root/ seeds/fruits/stem barks	Hypertension	E d'Avigdor, H Wohlmuth, Z Asfaw and T Awas [32], M Giday, T Teklehaymanot, A Animut and Y Mekonnen [35]
Root/ seeds/fruits/stem barks	Giardiasis	M Giday, T Teklehaymanot, A Animut and Y Mekonnen [35]
Root/ seeds/fruits/stem barks	Vomiting	S Zerabruk and G Yirga [23]

The primary method of preparation involves grinding the ingredients into a powder and mixing them with various components, such as coffee and traditional beverages. Subsequently, the final traditional medicine is administered orally, which is one of the most common routes for delivering traditional medicine. Several research findings indicated that the most commonly used routes of administration for herbal medicines are the oral route [25, 29, 37, 38], dermal [36] and nasal [39].

Furthermore, this species is also important for the treatments of diseases of livestock. For instance, the local community use a herbal medicine made from different parts of *C. aurea* to treat their livestock (Table 4). This implies that *C. aurea* is a versatile plant in its medicinal uses.

Table 3 Traditional use of *C. aurea* to treat diseases of livestock

Traditional used parts of <i>C.urena</i>	Locally used to treated	Reported by
Leaf /root/seeds/stem bark	Internal and external parasites	M Megersa, Z Asfaw, E Kelbessa, A Beyene and B Woldeab [22]; K Tolossa, E Debel, S

		Athanasiadou, A Tolera, G Ganga and JG Houdijk [30]; G Chekole, Z Asfaw and E Kelbessa [40]
Leaf /root/seeds/stem bark	Snakebite	M Megersa, Z Asfaw, E Kelbessa, A Beyene and B Woldeab [22];
Leaf /root/seeds/stem bark	External wound	B Adugna, NK GetachewTerefe, W Mamo and S Keskes [33]
Leaf /root/seeds/stem bark	Rabies disease	F Tamiru, W Terfa, E Kebede, G Dabessa, RK Roy and M Sorsa [41]
Leaf /root/seeds/stem bark	Cattle diarrhea/ cattle dysentery	H Tadege, E Mohammed, K Asres and T Gebre-Mariam [26]
Leaf	Eye disease	A Asfaw, E Lulekal, T Bekele, A Debella, E Debebe and B Sisay [42]

The preparation method involves chopping fresh or dried plant parts and mixing them with water, or using them without water. The application routes include topical [43, 44], nasal [45], and oral [46]. *C. aurea* can also serve as traditionally mosquito repellent in the area where the mosquito is prevalent [47]. Therefore, people who are living in mosquito prevalent areas can use it as malaria prevention mechanism. The whole part of the plant it is used as an insect repellent and its method of applications are laying the leaves of plants inside the house and rubbing onto clothes/skin and smoking the plant parts in the house [34, 38, 48, 49]. This implies that people can use *C. aurea* as an insect repellent, in addition to its applications in treating various ailments.

3.5 Photochemistry and pharmacological investigation reviews

The phytochemical analysis of extracts from medicinal plants is essential for discovering new phytochemical compounds, which plays a critical role in the development of novel drugs and alternative therapies.

Asaye Asfaw and Wondimagegn Mengist

Consequently, researchers worldwide have conducted studies on the phytochemical analysis and evaluation of the bioactive chemical constituents present in various parts of *C. aurea*. The result showed the presence of many secondary metabolites. For example, M Govindappa and M Poojashri [50] isolated alkaloids, terpenoids, saponins, cardiac glycosides, steroids, phenols, flavonoids, anthraquinone, and tannins from this distinguished species in South Africa. M Eyasu, W Shibeshi and M Giday [51], S Umer, A Tekewe and N Kebede [27], and Z Birhanu, M Wuhab and T Abula [52] also extracted the same bioactive compounds from the same parts of *C. aurea* in Ethiopia using similar methods. Moreover, compounds such as flavonols, proanthocyanidins, diterpenes, calpurnine, virgiline, virgiline pyrrolecarboxylic acid ester, and lapinine have also been identified in *C. aurea* [9, 43, 52]. This indicated that various types of secondary bioactive metabolites are present in *C. aurea*, which show promise for the development of various types new drugs, which might be solutions of the emergence of drug resistance bacteria.

In addition to ethnobotanical and phytochemical screening studies, numerous experimental research projects were carried out to assess the efficacy of secondary metabolites derived from various parts of *C. aurea* with the majority yielding encouraging outcomes. S Umer, A Tekewe and N Kebede [27] conducted an experimental investigation on the anti-bacterial effects of the bioactive chemicals screened from the leaf and stem of *C. aurea* and found that the extracts have strong anti-bacterial activities and showed promising effects for isolation and drug formulation. Similar studies [33, 43, 53] were also conducted on the anti-microbial effects of *C. aurea* extracts, and arrived at the same conclusion. The explanation for this is that wherever plant growth occurs, soil characteristics, rainfall, temperature, and other environmental factors do not influence the types of secondary bioactive metabolites.

Although a study by AA Adedapo, FO Jimoh, S Koduru, AJ Afolayan and PJ Masika [9] indicated that the antibacterial activities of *C. aurea* extracts are not as effective as standard drugs, the leaf extracts exhibit a broad spectrum of activity and demonstrate the highest efficacy against certain pathogenic bacterial species. However, their effectiveness is concentration-dependent. This indicates that various parts of *C. aurea* exhibit antimicrobial activities, although there are, some variations in the results that may be attributed to the concentration levels used, the timing of harvesting, and the types of pathogenic bacteria tested.

Additionally, research by M Eyasu, W Shibeshi and M Giday [51], B Adugna, NK GetachewTerefe, W Mamo

and S Keskes [33], and Z Birhanu, M Wuhab and T Abula [52] indicated that leaf extracts of *C. aurea* possess anti-protozoal properties, particularly against Plasmodium, amoeba, and Giardia species. This demonstrates that *C. aurea*, in addition to its antibacterial and antifungal activities, possesses antiparasitic properties against protozoa such as Plasmodium, amoebae, and Giardia species, which are prevalent in tropical countries, particularly in Ethiopia. The leaf extracts also exhibit anti-molluscicidal effects [54] and can also help lower blood pressure [32, 55]. This suggests that *C. aurea* not only has antimicrobial properties but also demonstrates anti-molluscicidal effects and contributes to the management of non-communicable diseases like hypertension.

In a study by A Zorloni, BL Penzhorn and JN Eloff [56], the leaf extracts were evaluated for their effectiveness against head lice and ticks, revealing significant and positive results in killing these pests. This suggests that *C. aurea* can also be utilized to deter external parasites. Furthermore, root extracts demonstrate anti-mycobacterial properties [26, 57], while extracts from the stem and bark show moderate anti-cancer effects [58]. This suggests that *C. aurea*, along with its microbial, anti-protozoal, anti-molluscicidal, and anti-external parasitic properties, may also be effective in treating severe illnesses such as cancer and tuberculosis, which are often difficult to manage with conventional medications.

4. Conclusion

This review indicated that *C. aurea* is a plant species traditionally used to treat various diseases in both humans and animals, particularly in different regions of Africa. Phytochemical analyses revealed that uses various parts—especially the leaves, stems, bark, and roots—contain secondary metabolites. While extracts from *C. aurea* have undergone experimental testing for various pharmacological properties, comprehensive scientific exploration and in-depth studies remain insufficient. Previous research has not adequately addressed the analysis of toxicity levels in bioactive secondary metabolites. Furthermore, results from in vitro experiments may not yield similar outcomes when applied under actual conditions. Although *C. aurea* has been used by local communities to treat a range of ailments—including rabies, snakebites, syphilis, diabetes mellitus, leishmaniasis, elephantiasis, tuberculosis, malaria, giardiasis, eye diseases in both cattle and humans, high blood pressure, amoebic dysentery, stomach pain, and cancer—there is currently no scientific evidence or in vitro testing to support its effectiveness for these conditions. Consequently, this gap necessitates further scientific investigation. Ethiopia is home to a diverse array of

Asaye Asfaw and Wondimagegn Mengist

potentially valuable medicinal plant species. However, factors such as changes in land use, excessive exploitation of natural resources, and overgrazing are contributing to a decline in both medicinal plants and the traditional practices associated with their uses. This situation underscores the need for a scientific approach to explore the applications of these plants. In conclusion, the authors recommend enhancing conservation efforts and thoroughly documenting traditional knowledge of *Calpurnia aurea* (Alt.) Benth and other potentially important medicinal plants.

References

1. Asfaw A, Lulekal E, Bekele T, Debella A, Abebe A, Degu S: Documentation of traditional medicinal plants use in Ensaro District, Ethiopia: implications for plant biodiversity and indigenous knowledge conservation. *Journal of Herbal Medicine* 2023, 38:100641.
2. Yedjou CG, Grigsby J, Mbemi A, Nelson D, Mildort B, Latinwo L, Tchounwou PB: The management of diabetes mellitus using medicinal plants and vitamins. *International journal of molecular sciences* 2023, 24(10):9085.
3. Lock J, MN F: *Legumes of the World*. London: Royal Botanic Gardens, Kew 2005.
4. Judd W, Campbell C, Kellog E, Stevens P, Donoghue M: *Plant Systematics: a phylogenetic approach*, Sinauer Axxoc, 287-292. In.: ISBN 0-87893-403-0; 2002.
5. Smýkal P, Coyne CJ, Ambrose MJ, Maxted N, Schaefer H, Blair MW, Berger J, Greene SL, Nelson MN, Besharat N: Legume crops phylogeny and genetic diversity for science and breeding. *Critical Reviews in Plant Sciences* 2015, 34(1-3):43-104.
6. Fathi G: Al-Sghair, Mohammed H. Mahklouf. Floristic Analysis and Species Diversity of the Family Fabaceae represented by voucher specimens Depending on the Flora of Libya. In., vol. 21: Species; 2020: 189-201.
7. Xu Z, Deng M, Xu Z, Deng M: Fabaceae or leguminosae. *Identification and Control of Common Weeds: Volume 2* 2017:547-615.
8. Duke J: *Handbook of legumes of world economic importance*: Springer Science & Business Media; 2012.
9. Adedapo AA, Jimoh FO, Koduru S, Afolayan AJ, Masika PJ: Antibacterial and antioxidant properties of the methanol extracts of the leaves and stems of *Calpurnia aurea*. *BMC Complementary and alternative medicine* 2008, 8(1):53.
10. Sita K, Sehgal A, HanumanthaRao B, Nair RM, Vara Prasad P, Kumar S, Gaur PM, Farooq M, Siddique KH, Varshney RK: Food legumes and rising temperatures: effects, adaptive functional mechanisms specific to reproductive growth stage and strategies to improve heat tolerance. *Frontiers in Plant Science* 2017, 8:1658.
11. Smith N: Fabaceae. In: *Amazon Fruits: An Ethnobotanical Journey*. edn.: Springer; 2023: 455-592.
12. Adenubi OT: Acaricidal efficacy of crude extracts and isolated flavonoids from *Calpurnia aurea* subsp. *aurea* against *Rhipicephalus turanicus*: University of Pretoria (South Africa); 2017.
13. Tebabal Kebede G: Phytochemical investigation, and assessment of antibacterial activities of *Calpurnia aurea* root bark. *Natural Products Chemistry Research*, 2021, 9:383.
14. Kenubih A, Belay M, Awol N, Birhan M: Extraction and phytochemical screening of *Calpurnia aurea* leaves and their effect against *malopaagus ovinus*. *Asian Journal of Medical and Pharmaceutical Researches* 2019, 9(1):1-10.
15. Nefo T, Dula T, Kelta B, Haile K: Isolation and Characterization of Flavonoid and Other Compounds from Seeds Extract of *Calpurnia Aurea*. *Extraction* 2019, 9(7).
16. Mulatu G: Antibacterial activities of *Calpurnia aurea* against selected animal pathogenic bacterial strains. *Advances in Pharmacological and Pharmaceutical Sciences* 2020, 2020(1):8840468.
17. Bogale A, Alemayehu H, Nedi T, Engidawork E: Antidiarrheal and antibacterial activities of *Calpurnia aurea*: Benth seed different extracts. *Evidence-based Complementary and Alternative Medicine* 2022, 2022(1):9582687.
18. Jansen PCM: *Spices, condiments and medicinal plants in Ethiopia, their taxonomy and agricultural significance*: [sn]; 1981.

Asaye Asfaw and Wondimagegn Mengist

19. Hedberg I: Flora of Ethiopia and Eritrea. In: The Biodiversity of African Plants. edn.: Springer; 1996: 802-804.
20. Wasihun Y, Alekaw Habteweld H, Dires Ayenew K: Antibacterial activity and phytochemical components of leaf extract of *Calpurnia aurea*. Scientific reports 2023, 13(1):9767.
21. Teklehaymanot T, Giday M: Ethnobotanical study of medicinal plants used by people in Zegie Peninsula, Northwestern Ethiopia. Journal of ethnobiology and Ethnomedicine 2007, 3(1):12.
22. Megersa M, Asfaw Z, Kelbessa E, Beyene A, Woldeab B: An ethnobotanical study of medicinal plants in Wayu Tuka district, east Welega zone of oromia regional state, West Ethiopia. Journal of ethnobiology and ethnomedicine 2013, 9(1):68.
23. Zerabruk S, Yirga G: Traditional knowledge of medicinal plants in Gindeberet district, Western Ethiopia. South African journal of botany 2012, 78:165-169.
24. Kidane B, van Andel T, van der Maesen LJG, Asfaw Z: Use and management of traditional medicinal plants by Maale and Ari ethnic communities in southern Ethiopia. Journal of ethnobiology and ethnomedicine 2014, 10(1):46.
25. Teklehaymanot T: Ethnobotanical study of knowledge and medicinal plants use by the people in Dek Island in Ethiopia. Journal of Ethnopharmacology 2009, 124(1):69-78.
26. Tadeg H, Mohammed E, Asres K, Gebre-Mariam T: Antimicrobial activities of some selected traditional Ethiopian medicinal plants used in the treatment of skin disorders. Journal of ethnopharmacology 2005, 100(1-2):168-175.
27. Umer S, Tekewe A, Kebede N: Antidiarrhoeal and antimicrobial activity of *Calpurnia aurea* leaf extract. BMC complementary and alternative medicine 2013, 13(1):21.
28. Alemayehu G, Asfaw Z, Kelbessa E: Ethnobotanical study of medicinal plants used by local communities of Minjar-Shenkora District, North Shewa Zone of Amhara Region, Ethiopia. Journal of Medicinal Plants Studies 2015, 3(6):1-11.
29. Regassa R, Bekele T, Megersa M: Ethnobotanical study of traditional medicinal plants used to treat human ailments by halaba people, southern Ethiopia. Journal of Medicinal Plants 2017, 5(4):36-47.
30. Tolossa K, Debela E, Athanasiadou S, Tolera A, Ganga G, Houdijk JG: Ethno-medicinal study of plants used for treatment of human and livestock ailments by traditional healers in South Omo, Southern Ethiopia. Journal of Ethnobiology and Ethnomedicine 2013, 9(1):32.
31. Waka M, Hopkins RJ, Curtis C: Ethnobotanical survey and testing of plants traditionally used against hematophagous insects in Eritrea. Journal of ethnopharmacology 2004, 95(1):95-101.
32. d'Avigdor E, Wohlmuth H, Asfaw Z, Awas T: The current status of knowledge of herbal medicine and medicinal plants in Fiche, Ethiopia. Journal of ethnobiology and ethnomedicine 2014, 10(1):38.
33. Adugna B, GetachewTerefe NK, Mamo W, Keskes S: Potential In vitro Anti-Bacterial Action of Selected Medicinal Plants Against *Escherichia coli* and Three *Salmonella* Species. International Journal, 2014, 5(2):85-89.
34. Yineger H, Yewhalaw D, Teketay D: Ethnomedicinal plant knowledge and practice of the Oromo ethnic group in southwestern Ethiopia. Journal of Ethnobiology and Ethnomedicine 2008, 4(1):11.
35. Giday M, Teklehaymanot T, Animut A, Mekonnen Y: Medicinal plants of the Shinasha, Agew-awi and Amhara peoples in northwest Ethiopia. Journal of Ethnopharmacology 2007, 110(3):516-525.
36. Agisho H, Osie M, Lambore T: Traditional medicinal plants utilization, management and threats in Hadiya Zone, Ethiopia. Journal of Medicinal Plant Studies 2014, 2(2):94-108.
37. Birhanu Z: Ethno-botanical survey on medicinal plants used by ethnic groups of Denbia district, north-western Ethiopia. Journal of Natural Remedies 2011, 11(2):119-123.
38. Kidane D, Tomass Z, Dejene T: Community knowledge of traditional mosquito repellent plants in Kolla Temben District, Tigray,

Asaye Asfaw and Wondimagegn Mengist

- Northern Ethiopia. Scientific Research and Essays 2013, 8(24):1139-1144.
39. Asnake S, Teklehaymanot T, Hymete A, Erko B, Giday M: Survey of medicinal plants used to treat malaria by Sidama People of Boricha District, Sidama Zone, South Region of Ethiopia. Evidence-Based Complementary and Alternative Medicine 2016, 2016.
 40. Chekole G, Asfaw Z, Kelbessa E: Ethnobotanical study of medicinal plants in the environs of Tara-gedam and Amba remnant forests of Libo Kemkem District, northwest Ethiopia. Journal of ethnobiology and ethnomedicine 2015, 11(1):4.
 41. Tamiru F, Terfa W, Kebede E, Dabessa G, Roy RK, Sorsa M: Ethnoknowledge of plants used in veterinary practices in Dabo Hana District, West Ethiopia. Journal of Medicinal plants Research 2013, 7(40):2960-2971.
 42. Asfaw A, Lulekal E, Bekele T, Debella A, Debebe E, Sisay B: Medicinal plants used to treat livestock ailments in Ensaro district, North Shewa Zone, Amhara regional state, Ethiopia. BMC Veterinary Research 2022, 18(1):235.
 43. Soyelu O, Masika P: Antibacterial and antioxidant activities of some selected plants used for the treatment of cattle wounds in the Eastern Cape. African Journal of Biotechnology 2012, 11(14):3476-3481.
 44. Teklay A, Abera B, Giday M: An ethnobotanical study of medicinal plants used in Kilte Awulaelo District, Tigray Region of Ethiopia. Journal of ethnobiology and ethnomedicine 2013, 9(1):65.
 45. Yigezu Y, Haile DB, Ayen WY: Ethnoveterinary medicines in four districts of Jimma zone, Ethiopia: cross sectional survey for plant species and mode of use. BMC veterinary research 2014, 10(1):76.
 46. Bekalo TH, Woodmatas SD, Woldemariam ZA: An ethnobotanical study of medicinal plants used by local people in the lowlands of Konta Special Woreda, southern nations, nationalities and peoples regional state, Ethiopia. Journal of Ethnobiology and Ethnomedicine 2009, 5(1):26.
 47. Hiruy B, Getu E: Evaluation of *Calpurnia aurea* leaf extracts as natural insect repellents for stored product insect pests in Ethiopia. All Life 2023, 16(1):2163303.
 48. Yirga G, Teferi M, Gidey G, Zerabruk S: An ethnoveterinary survey of medicinal plants used to treat livestock diseases in Seharti-Samre district, Northern Ethiopia. African Journal of Plant Science 2012, 6(3):113-119.
 49. Bekele D, Asfaw Z, Petros B, Tekie H: Ethnobotanical study of plants used for protection against insect bite and for the treatment of livestock health problems in rural areas of Akaki District, Eastern Shewa, Ethiopia. Topclass Journal of Herbal Medicine 2012, 1(2):12-24.
 50. Govindappa M, Poojashri M: Antimicrobial, antioxidant and in vitro anti-inflammatory activity of ethanol extract and active phytochemical screening of *Wedelia trilobata* (L.) Hitchc. Journal of Pharmacognosy and Phytotherapy 2011, 3(3):43-51.
 51. Eyasu M, Shibeshi W, Giday M: In vivo antimalarial activity of hydromethanolic leaf extract of *Calpurnia aurea* (Fabaceae) in Mice infected with chloroquine sensitive *Plasmodium berghei*. International Journal of Pharmacy and Pharmacology 2013, 2:131-142.
 52. Birhanu Z, Wuhab M, Abula T: Antimalarial activity of *calpurnia aurea* hydroalcoholic leaf extract in mice infected with *Plasmodium berghei*. Pharmacologyonline 2015, 2:73-79.
 53. Muk L, Naidoo V, Eloff JN: In vitro antibacterial activity of seven plants used traditionally to treat wound myiasis in animals in Southern Africa. Journal of Medicinal Plants Research 2012, 6(27):4379-4388.
 54. Kubo I, Matsumoto T, Kozuka M, Chapya A, Naoki H: Quinolizine alkaloids from the African medicinal plant *Calpurnia aurea*: molluscicidal activity and structural Study by 2D-NMR. Agricultural and biological chemistry 1984, 48(11):2839-2841.
 55. Getiye Y, Tolessa T, Engidawork E: Antihypertensive activity of 80% methanol seed extract of *Calpurnia aurea* (Ait.) Benth. subsp. *aurea* (Fabaceae) is mediated through calcium antagonism induced vasodilation. Journal of ethnopharmacology 2016, 189:99-106.
 56. Zorloni A, Penzhorn BL, Eloff JN: Extracts of *Calpurnia aurea* leaves from southern Ethiopia

Asaye Asfaw and Wondimagegn Mengist

attract and immobilise or kill ticks. *Veterinary parasitology* 2010, 168(1-2):160-164.

57. Gemechu A, Giday M, Worku A, Ameni G: In vitro anti-mycobacterial activity of selected medicinal plants against *Mycobacterium tuberculosis* and *Mycobacterium bovis* strains. *BMC complementary and alternative medicine* 2013, 13(1):291.
58. Korir E, Kiplimo JJ, Crouch NR, Moodley N, Koorbanally NA: Isoflavones from *Calpurnia aurea* subsp. *aurea* and their anticancer activity. *African Journal of Traditional, Complementary and Alternative Medicines* 2014, 11(5):33-37.